

Draft Guidance on Prochlorperazine Maleate

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

Active Ingredient: Prochlorperazine maleate

Dosage Form; Route: Tablet; oral

Recommended Studies: Two studies

1. Type of study: Fasting
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: EQ 10 mg Base
Subjects: Males and non-pregnant, non-lactating females, general population
Additional comments: None

2. Type of study: Fed
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: EQ 10 mg Base
Subjects: Males and non-pregnant, non-lactating females, general population
Additional comments: None

Analyte to measure: Prochlorperazine in plasma

Bioequivalence based on (90% CI): Prochlorperazine

Waiver request of in vivo testing: EQ 5 mg Base mg based on (i) acceptable bioequivalence studies on the EQ 10 mg Base strength, (ii) acceptable in vitro dissolution testing of both strengths, and (iii) proportional similarity of the formulations between both strengths

Dissolution test method and sampling times: The dissolution information for this drug product can be found in the FDA's Dissolution Methods database, <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Conduct comparative dissolution testing on 12 dosage units each of both strengths of the test and reference products. Specifications will be determined upon review of the abbreviated new drug application.