

Contains Nonbinding Recommendations
Draft - Not for Implementation
Draft Guidance on Naproxen Sodium
February 2023

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA's guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency's current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredient: Naproxen sodium

Dosage Form; Route: Tablet, extended release; oral

Recommended Studies: Two in vivo bioequivalence studies with pharmacokinetic endpoints

1. Type of study: Fasting
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: EQ 750 mg Base
Subjects: Healthy males and non-pregnant, non-lactating females
Additional comment: None

2. Type of study: Fed
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: EQ 750 mg Base
Subjects: Healthy males and non-pregnant, non-lactating females
Additional comment: None

Analyte to measure: Naproxen in plasma

Bioequivalence based on (90% CI): Naproxen

Ensure the time to maximum observed plasma concentration (T_{max}) from the test and reference products support comparable therapeutic outcome.

Additional strengths: Bioequivalence of the EQ 375 mg Base and EQ 500 mg Base strengths to the corresponding reference product strengths may be demonstrated based on principles laid out in the most recent version of the FDA guidance for industry on *Bioequivalence Studies with Pharmacokinetic Endpoints for Drugs Submitted Under an Abbreviated New Drug Application*.^a

Dissolution test method and sampling times: For modified release drug products, applicants should develop specific discriminating dissolution methods. Alternatively, applicants may use the dissolution method set forth in any related official United States Pharmacopeia (USP) drug product monograph, or in the FDA's Dissolution Methods database, <http://www.accessdata.fda.gov/scripts/cder/dissolution/>, provided that applicants submit adequate dissolution data supporting the discriminating ability of such a method. If a new dissolution method is developed, submit the dissolution method development and validation report with the complete information/data supporting the proposed method. Conduct comparative dissolution testing on 12 dosage units for each strength of the test and reference products. Specifications will be determined upon review of the Abbreviated New Drug Application (ANDA).

In addition to the method above, submit dissolution profiles on 12 dosage units for each strength of the test and reference products generated using USP Apparatus 1 at 100 rpm and/or Apparatus 2 at 50 rpm in at least three dissolution media (e.g., pH 1.2, 4.5 and 6.8 buffer). Agitation speeds may be increased if appropriate. It is acceptable to add a small amount of surfactant if necessary. Include early sampling times of 1, 2, and 4 hours and continue every 2 hours until at least 80% of the drug is released to provide assurance against premature release of drug (dose dumping) from the formulation.

Revision History: Recommended April 2010; Revised February 2023

Unique Agency Identifier: PSG_020353

^a For the most recent version of a guidance, check the FDA guidance web page at <https://www.fda.gov/regulatory-information/search-fda-guidance-documents>.